

Foliar fungicides registered for use on barley in WA – 2024-25

This information is a guide only. Chemical labels should be read, and rates should be checked before using fungicides. Mention of trade names does not imply endorsement of any company's products. Rates of registered products in mL/ha or g/ha. Symbols used refer to example product trade names.

Table 1 Fungicide active ingredient and equivalent trade names registered in barley. List may not be comprehensive

Active ingredient	Fungicide group (FRAC code)	Leaf rust	Stem rust	Scald	Spot form net blotch	Net form net blotch	Powdery mildew	Example product trade names (® = registered trademark; other symbols refer to rates in the table for that specific product).	Withholding period (harvesting or grazing, in weeks)
Azoxystrobin 200g/L + Cyproconazole 80g/L	11 + 3	200-800	No registration	No registration Amistar® Xtra only 400-800	200-800	200-800	400-800	Amazer® 280, Amistar® Xtra, Azoxy® Xtra, Azoxy® Cypro, Azoxyguard® Xtra, Azoxystrobin® Extra, Roxy® Extra	Harvest-6, Grazing-3
Azoxystrobin 75g/L + Epoxiconazole 75g/L	11 + 3	420-840	No registration	420-840	420-840 Radial® only	420-840	420-840	Radial® (Eyespot suppression at 840mL), Serial® 150 EC	Harvest-6, Grazing-6
Azoxystrobin 80g/L + Epoxiconazole 31.25g/L	11 + 3	500-2000	1000-2000	1000-2000	1000-2000	1000-2000	1000-2000	Tazer® Xpert	Harvest-N/A, Grazing-3

Active ingredient	Fungicide group (FRAC code)	Leaf rust	Stem rust	Scald	Spot form net blotch	Net form net blotch	Powdery mildew	Example product trade names (® = registered trademark; other symbols refer to rates in the table for that specific product).	Withholding period (harvesting or grazing, in weeks)
Azoxystrobin 120g/L + Epoxiconazole 100g/L	11 + 3	313-625	No registration	313	313-625	313-625	313	Avior Gold	Harvest - 6, Grazing -6
Azoxystrobin 320g/L + Epoxiconazole 250g/L	11 + 3	125 – 250		125 - 250	125 – 250	125 - 250	125 – 250	Radial® Opti (Eyespot suppression at 250mL)	Harvest – 0 when used as directed. Grazing - 3
Azoxystrobin 120g/L + Tebuconazole 200g/L	11 + 3	315- 630	No registration	315	315- Suppression only	315- 630 Suppression only	315-630	Veritas®, Azlan®, Stunner®320,	Harvest-6, Grazing-3
Azoxystrobin 133g/L + Prothioconazole 100g/L	11 + 3	300 - 600	No registration	300 - 600	300 - 600	300 - 600	300 - 600	Maxentis®EC **Also registered for Ramularia Leaf Spot	Harvest-5, Grazing-3
Azoxystrobin 222g/L + Tebuconazole 370g/L	11 + 3	170-340	No registration	170	170-340	170-340	170-340	Veritas Opti®	Harvest-6, Grazing-3
Azoxystrobin 200g/L + Flutriafol 125g/L	11 + 3	No registration	No registration	No registration	800	No registration	No registration	Metropol®SC	Harvest-10, Grazing-10
Azoxystrobin 250g/L + Epoxiconazole 125g/L + Tebuconazole 125g/L	11 + 3 + 3	No registration	No registration	250	No registration	250	No registration	Triaxe® Fungicide	Harvest-6, Grazing-6
Azoxystrobin 200g/L + Propiconazole 200g/L	11 + 3	300-600	No registration	300-600	300-600	300-600	200-600	Topnotch® - also registered at 600mL for suppression of eyespot & physiological leaf spot	Harvest-4, Grazing-3
Azoxystrobin 280g/L + Prothioconazole 120g/L	11 + 3	263 - 525	No registration	No registration	263 - 525	263 - 525	263 - 525	Uplift®	Harvest-6, Grazing-3
Azoxystrobin 600g/kg + Cyproconazole 240g/kg	11 + 3	67-267g	No registration	No registration	67-267g	67-267g	133-267g	Azoxycypro® WG, Kingstar®	Harvest-6, Grazing-3

Active ingredient	Fungicide group (FRAC code)	Leaf rust	Stem rust	Scald	Spot form net blotch	Net form net blotch	Powdery mildew	Example product trade names (® = registered trademark; other symbols refer to rates in the table for that specific product).	Withholding period (harvesting or grazing, in weeks)
Epoxiconazole 125g/L	3	250–500 ¹ & ²	No registration	250 ¹	No registration	250–500 ¹ Fungicide resistance warning (see footer)	250 ¹	1: Opus®125, Soprano®, Rivet® 125, Epoxiconazole 125 (4Farmers/Accensi brands only), Epoxiguard®125, Bachlor® 125SC, Epoxi® 125SC, Epoxicon®125 SC, Epoxy® 125 & SC, Overture®, Protector 125, Epoxistar® 125SC, Tangent® 125SC 2: Epoch®125 & SC, Epoxiconazole 125 & 125SC (all other brandnames), Swoosh® 125, Vindicate® 125SC, Epoxikey®	Harvest-6, Grazing-6
Epoxiconazole 500g/L	3	65-125 ³ 62-125 ⁴	No registration	65-125 ³ 62.5 ⁴	65-125 ³ 62.5-125 ⁴ Fungicide resistance warning (see footer)	65-125 ³ 62.5-125 ⁴ Fungicide resistance warning (see footer)	65-125 ³ 62.5 ⁴	3. Soprano® 500, Epoxiconazole 500 (<i>All registered for Eyespot suppression 125mL</i>) 4. Epoxiconazole 500 WDG	3. Harvest-not required when used as directed, Grazing-3 4. Harvest-6, Grazing-6
Epoxiconazole 750g/kg	3	42-84	No registration	42	No registration	42-84 Fungicide resistance warning (see footer)	42	Epoxy® 750 WG	Harvest-6, Grazing-6
Epoxiconazole 800g/kg	3	39-78	No registration	No registration	No registration	No registration	No registration	Octopus® 800 WG	Harvest-6, Grazing-6
Epoxiconazole 62.5g/L+ Pyraclostrobin 85g/L	3 + 11	500	No registration	500	500-1000	500-1000 Fungicide resistance warning (see footer)	500-1000	Opera®, Pirastar®	Harvest-0 if not applied after Z59, Grazing-3

Active ingredient	Fungicide group (FRAC code)	Leaf rust	Stem rust	Scald	Spot form net blotch	Net form net blotch	Powdery mildew	Example product trade names (® = registered trademark; other symbols refer to rates in the table for that specific product).	Withholding period (harvesting or grazing, in weeks)
Flutriafol 250g/L	3	No registration	No registration	No registration	No registration	No registration	250-500 Fungicide resistance warning (see footer)	Bayonet®, Flutifol® 250, Flutriafol 250 & SC, Flufo®, Impact®, Tricam®, Intake® Pro, Pollux®, Flout® 250SC, Futura® 250SC, Flukey®	Harvest-10, Grazing-10
Flutriafol 500g/L	3	No registration	No registration	No registration	No registration	No registration	125–250 Fungicide resistance warning (see footer)	Impact® Endure, Jubilee®500 & Loaded, Leda® 500, Jubilee® 500, Flutriafol MAX, ****Flutriafol 500 and 500SC only certain brands registered as foliar spray	Harvest-10, Grazing-10
Mefentrifluconazole 100g/L + Fluxapyroxad 50g/L	3+7	750	750	750	750	750	750	Revystar®	Harvest-0 when used as directed, Grazing-4
Fluxapyroxad 62.5g/L	7	250 or 500-1000	No registration	250 or 500-1000	250 or 500-1000 Fungicide resistance warning (see footer)	250 or 500-1000	500-1000	Imbrex® Fungicide	Harvest-0 if not applied after Z59, Grazing-2
Epoxiconazole 42g/L + Fluxapyroxad 42g/L + Pyraclostrobin 67g/L	3 + 7 + 11	750	No registration	750	750	750	750	Cerix®	Harvest-not required when used as directed, Grazing-6

Active ingredient	Fungicide group (FRAC code)	Leaf rust	Stem rust	Scald	Spot form net blotch	Net form net blotch	Powdery mildew	Example product trade names (® = registered trademark; other symbols refer to rates in the table for that specific product).	Withholding period (harvesting or grazing, in weeks)
Propiconazole 250g/L	3	No registration	No registration	500 ^{5 & 6} Or 150–500 ⁷ Rate varies for Propi conazole 250 brands – check labels	500 ^{5 & 8} or 250-500 ⁶ Or 150–500 ⁷ Fungicide resistance warning (see footer)	250–500 ⁶ (Tilt® and Genfarm Propiconazole 250 only) Fungicide resistance warning (see footer)	150–500 * *Generally all products but check labels. Fungicide resistance warning (see footer)	5: Cracker®250EC, Prestige®, Detour®250EC, Procon®250EC, Propi® 250, Propicol®250EC, Propiconazole 250 & 250EC (all other brands), Petulant® 250EC, Pace®250EC, Restore®250EC, Slipstream®250EC, Colossus 250 6: Tilt® 250, Propiconazole 250EC (Genfarm) 7: Propiconazole 250 (Titan brands only). Propicon 250EC, 8: Procure (Only for Spot form net blotch)	Harvest-4, Grazing-1
Propiconazole 250g/L + Benzovindiflupyr 40g/L	3 + 7	500	No registration	500	500	500	500	Elatus Ace® * Also registered for Ramularia leaf spot and Ring spot *	Harvest-0 if used as directed, Grazing-1.5
Propiconazole 435g/L	3	No registration	No registration	285	285 Fungicide resistance warning (see footer)	No registration Fungicide resistance warning (see footer)	85–285 Fungicide resistance warning (see footer)	Fitness® Fungicide	Harvest-4, Grazing-1

Active ingredient	Fungicide group (FRAC code)	Leaf rust	Stem rust	Scald	Spot form net blotch	Net form net blotch	Powdery mildew	Example product trade names (® = registered trademark; other symbols refer to rates in the table for that specific product).	Withholding period (harvesting or grazing, in weeks)
Propiconazole 500g/L	3	125-250 NOT Propiconazole500 or Propi Super	No registration	250	125-250 Fungicide resistance warning (see footer)	125-250 NOT Propiconazole500 or Propi Super Fungicide resistance warning (see footer)	75-250 Fungicide resistance warning (see footer)	Throttle® 500, Prop® 500, Tilt® 500EC, Propiconazole 500EC, Propi Super 500	Harvest-4, Grazing-1
Propiconazole 550g/L	3	No registration	No registration	230	230 Fungicide resistance warning (see footer)	No registration	70-230 Fungicide resistance warning (see footer)	Cracker Jack®550EC, Propiconazole 550EC, Pacer® 550EC, Prestige® 550, Picaro Hi-Load 550EC, Propicon® 550EC	Harvest-4, Grazing-1
Propiconazole 625g/L	3	No registration	No registration	200	200 Fungicide resistance warning (see footer)	No registration	60-200 Fungicide resistance warning (see footer)	Bumper® 625EC, Propiconazole 625EC, Pinazole®	Harvest-4, Grazing-1
Propiconazole 750g/L	3	No registration	No registration	170	170 Fungicide resistance warning (see footer)	No registration	50-170 Fungicide resistance warning (see footer)	Cracker Jack® 750EC	Harvest-4, Grazing-1
Propiconazole 800g/L	3	No registration	No registration	160	160	No registration	45-160	Cracker Jack® 800EC	Harvest-4, Grazing-1

Active ingredient	Fungicide group (FRAC code)	Leaf rust	Stem rust	Scald	Spot form net blotch	Net form net blotch	Powdery mildew	Example product trade names (® = registered trademark; other symbols refer to rates in the table for that specific product).	Withholding period (harvesting or grazing, in weeks)
Propiconazole 250g/L + Cyproconazole 80g/L	3 + 3	250-500	No registration	500	250 -500	250 - 500	150-500 Fungicide resistance warning (see footer)	Tilt® Xtra	Harvest-6, Grazing-3
Propiconazole 250g/L + Tebuconazole 250g/L	3 + 3	125-250	No registration	250	125-250 Fungicide resistance warning (see footer)	125-250 Fungicide resistance warning (see footer)	125-250 Fungicide resistance warning (see footer)	Cogito® (Discontinued)	Harvest-5, Grazing-2
Prothioconazole 150g/L + Bixafen 75g/L	3 + 7	400-500	No registration	300-500	300-500	300-500	300-500	Aviator Xpro®	Harvest –NA if used as directed, Grazing 4
Prothioconazole 250g/L	3	250 ⁹	No registration	250 ^{9 10}	250 ¹⁰	250 ⁹ Or 125-250 ¹⁰	250 ^{9 10}	9: Protek® 250EC 10: Proviso® If high disease pressure- see tank mixes	Harvest- 5, Grazing-2
Prothioconazole 480g/L	3	70-130	No registration	70-130	70-130	70-160	70-130	Proline® 480 * Also registered for physiological leaf spotting 70–130 *	Harvest- 5, Grazing-2
Quinoxifen 250g/L	13	No registration	No registration	No registration	No registration	No registration	200-300	Legend®	Harvest-not required when used as directed, Grazing-7
Spiroxamine 500g/L	5	No registration	No registration	No registration	No registration	No registration	600	Prosper®500EC	Harvest-not required when used as directed, Grazing-6

Active ingredient	Fungicide group (FRAC code)	Leaf rust	Stem rust	Scald	Spot form net blotch	Net form net blotch	Powdery mildew	Example product trade names (® = registered trademark; other symbols refer to rates in the table for that specific product).	Withholding period (harvesting or grazing, in weeks)
Tebuconazole 430g/L	3	No registration	No registration	145	No registration	No registration	145-290 ¹ Fungicide resistance warning (see footer)	Laguna®430, Orius®430SC, Rebuke®430SC, Zolo® 430SC, Stingray®, Tebucol®430, Tebucon®430, Whirlwind®430, Toledo 430SC, Tebazal®430, Tebuconazole, 430 & SC, Tripod®, Cure® 430, Tusk®, Tebby® 430, Maya Tebu® 430, Tebustar® 430SC,	Harvest-5, Grazing-2
Tebuconazole 750g/kg	3	No registration	No registration	83	No registration	No registration	83-166 Fungicide resistance warning (see footer)	Tebuconazole 750WDG, Buzz® Ultra 750 WG, Ultrateb® 750 WG	Harvest-5, Grazing-2
Tebuconazole 800g/kg	3	No registration	No registration	78	No registration	No registration	78-156 Fungicide resistance warning (see footer)	Tebuconazole 800WG, Turbulence® 800WG, Laguna Xtreme®800 WG	Harvest-5, Grazing-2
Tebuconazole 210g/L + Prothioconazole 210g/L	3 + 3	150-300	No registration	150-300	150-300 Fungicide resistance warning (see footer)	150-300 Fungicide resistance warning (see footer)	150-300	Prosaro®420SC, Mantaray® 420SC, Mightyazole® 420, Pro Grow® 420SC, Progress 420EC, Prothioconazole® T, Pro Teb® SC & 420SC, Prothio T® & 420, Prothioconazole and Tebuconazol, Teb Pro®420, Tebupro®420SC	Harvest-5, Grazing-2

Active ingredient	Fungicide group (FRAC code)	Leaf rust	Stem rust	Scald	Spot form net blotch	Net form net blotch	Powdery mildew	Example product trade names (® = registered trademark; other symbols refer to rates in the table for that specific product).	Withholding period (harvesting or grazing, in weeks)
Tebuconazole 300g/L + Prothioconazole 300g/L	3 + 3	105-210	No registration	105-210	105-210 Fungicide resistance warning (see footer)	105-210 Fungicide resistance warning (see footer)	105-210	Coprotec® Xtra	Harvest-5, Grazing-2
Tebuconazole 45g/kg + Sulphur 700g/kg	3 + M2	No registration	No registration	1370	No registration	No registration	1370-2750	Unicorn® 745WG (for feed barley)	Harvest-5, Grazing-2
Triadimefon 125g/L	3	No registration	No registration	No registration	No registration	No registration	1000 Fungicide resistance warning (see footer)	Triadimefon 125EC, Slingshot®	Harvest-5, Grazing-2
Triadimefon 500g/L	3	No registration	No registration	No registration	No registration	No registration	250 Fungicide resistance warning (see footer)	Triadimefon 500 WP / WG	Harvest-4 Grazing- not to be sold for animal consumption

Table 2 Tank mixes registered in barley. Refer to labels for application rates, guidelines on application and crop safety.

Active ingredient + product to mix with	Fungicide groups (FRAC codes)	Leaf rust	Stem rust	Scald	Spot form net blotch	Net form net blotch	Powdery mildew	Example product trade names (@ = registered trademark; other symbols refer to rates in the table for that specific product).	Withholding period (harvesting or grazing, in weeks)
Prothioconazole 250g/L + Orius®	3 + 3	125 +75 Low disease 250 + 145 Mod – high disease	No registration	250 Proviso only or 125 + 75 Low disease 250 + 145 Mod – High disease	250 Proviso only or 125 + 75 Low disease 250 + 145 Mod – High disease Fungicide resistance warning (see footer)	125-250 Proviso only Low – Mod disease 125-250 + 75-145 Mod – high disease Fungicide resistance warning (see footer)	250 Proviso only or 125 + 75 Low disease 250 + 145 Mod – High disease	Proviso® * Also registered for low to moderate risk Physiological Leaf Spot 125-250 + 75 *	Harvest-5, Grazing-2
Prothioconazole 250g/L + Veritas®	3 + (3 + 11)	250 + 315 Mod – High disease	No registration	250 + 315 Mod – High disease	250 + 315 Mod – High disease	250 + 315 High disease	250 + 315 Mod – High disease	Proviso® * Also registered for high-risk Physiological Leaf Spot 250 + 315 *	Harvest-5, Grazing-2
Prothioconazole 250g/L + Laguna® (Tebuconazole 430g/L)	3 + 3	125-250 + 75-150	No registration	125-250 + 75-150	125-250 + 75-150 Fungicide resistance warning (see footer)	125-250 + 75-150 Fungicide resistance warning (see footer)	125-250 + 75-150	Protek® 250EC	Harvest-5, Grazing-2
Prothioconazole 250g/L + Overture® (Epoconazole 125g/L)	3 + 3	250 + 250 or 500	No registration	250 + 250 or 500	No registration	250 + 250 or 500	250 + 250 or 500	Protek® 250EC	Harvest-5, Grazing-2

Active ingredient + product to mix with	Fungicide groups (FRAC codes)	Leaf rust	Stem rust	Scald	Spot form net blotch	Net form net blotch	Powdery mildew	Example product trade names (® = registered trademark; other symbols refer to rates in the table for that specific product).	Withholding period (harvesting or grazing, in weeks)
Prothioconazole 250g/L + Accolade® (Azoxystrobin 250g/L)	3 + 11	250 + 320 or 640	No registration	No registration	250 + 320 or 640	250 + 320 or 640	No registration	Protek® 250EC	Harvest-5, Grazing-2
Prothioconazole 420g/L + Tebuconazole 430 SC	3 + 3	75-150 + 74-148	No registration	75-150 + 74-148	75-150 + 74-148	75-150 + 74-148	75-150 + 74-148	Genfarm Prothioconazole 420 SC, Relyon Prothioconazole 420 SC. Need to include adjuvant	Harvest-4, Grazing-4
Prothioconazole 480g/L + Conquest Stingray®430	3 + 3	66-132 + 74-148	No registration	66-132 + 74-148	66-132 + 74-148 Fungicide resistance warning (see footer)	66-132 + 74-148 Fungicide resistance warning (see footer)	66-132 + 74-148	Stallion® 480EC	Harvest-4, Grazing-4
Prothioconazole 480g/L + Orius® 430 SC	3 + 3	70 – 130 + 75 - 145	No registration	70 – 130 + 75 - 145	70 – 130 + 75 - 145	No registration	70 – 130 + 75 - 145	Proline® 480 Also registered for physiological leaf spotting 70 – 130 + 75 - 145	Harvest- 5, Grazing-2
Prothioconazole 800g/kg + Imtrade Turbulence® 800WG	3 + 3	40-80 + 40-80	No registration	40-80 + 40-80	40-80 + 40-80 Fungicide resistance warning (see footer)	40-80 + 40-80 Fungicide resistance warning (see footer)	40-80 + 40-80	Dalbie® 800WG	Harvest-5, Grazing-2
Azoxystrobin 250g/L + Overture® (or equivalent Epoxiconazole 125g/L)	11 + 3	320-640 + Label rate	No registration	320-640 + Label rate	No registration	320-640 + Label rate	320-640 + Label rate	Accolade® 250EC	Harvest-6, Grazing-3

Active ingredient + product to mix with	Fungicide groups (FRAC codes)	Leaf rust	Stem rust	Scald	Spot form net blotch	Net form net blotch	Powdery mildew	Example product trade names (@ = registered trademark; other symbols refer to rates in the table for that specific product).	Withholding period (harvesting or grazing, in weeks)
Azoxystrobin 250g/L + Laguna® (or equivalent Tebuconazole 430g/L)	11 + 3	No registration	No registration	160 + Label rate	No registration	No registration	160-320 + Label rate	Accolade® 250EC	Harvest-6, Grazing-3
Azoxystrobin 625g/L + Soprano® 500	11 + 3	65-250 + 65-125	No registration	65-250 + 65-125	65-250 + 65-125	65-250 + 65-125	65-250 + 65-125	Mirador®625	Harvest-4, Grazing-3
Azoxystrobin 625g/L + Orius® 430 SC	11 + 3	65-125 + 145-290	No registration	65-125 + 145-290	65-125 + 145-290	65-125 + 145-290	65-125 + 145-290	Mirador®625	Harvest-6, Grazing-3
Azoxystrobin 625g/L + Bumper®625	11 + 3	100-200 + 100-200	No registration	100-200 + 100-200	100-200 + 100-200	100-200 + 100-200	100-200 + 100-200	Mirador®625	Harvest-4, Grazing-3
Azoxystrobin 625g/L + Proviso®	11 + 3	65-250 + 125-250	No registration	65-250 + 125-250	65-250 + 125-250	65-250 + 125-250	65-250 + 125-250	Mirador®625 * Also registered for Ramularia leaf spot *	Harvest-5, Grazing-3
Pyraclostrobin 250g/L + Epoxiconazole 125g/L	11 + 3	170 + 250	No registration	170 + 250	170-340 + 250-500	170-340 + 250-500	170-340 + 250-500	Pyraclostrobin 250EC	Harvest-Not required when used as directed, Grazing-3

Notes: Procure® is registered only for spot form net blotch in barley and none of the other diseases.

Carefully read labels of fungicides with active ingredient Epoxiconazole 125g/L, as registrations vary; some are not registered for scald, powdery mildew, or net-form net blotch.

Carefully read labels of fungicides with active ingredient Propiconazole 250g/L, as registrations and rates vary; many are not registered for leaf rust, or net-form net blotch.

Fungicide resistance warning

Fungicide resistance to some groups of fungicides compromises control of barley powdery mildew, spot-form net blotch, and net-form net blotch. These fungicides, marked in table with a warning, provide variable and often poor levels of control and may promote further mutations of

the fungicide resistant strains of each disease. They are not recommended and, if possible, you should use another registered fungicide instead. Fungicides in other fungicide groups are registered to manage these diseases in WA.

Where possible it is best to use the currently effective fungicide actives only once per season to reduce selection pressure on the powdery mildew fungus. For greatest efficacy, fungicides should be applied before significant levels of disease establish in crop - use the fungicide to protect the crop rather than attempt to recover it. Fungicide resistance has come about from continuous use of fungicides with same actives or from same group as seed dressings and following up with multiple foliar applications.

Barley powdery mildew: Resistance and reduced sensitivity to Group 3 fungicides (tebuconazole, propiconazole, flutriafol) has been confirmed in WA.

Barley net-form net blotch: Resistance and reduced sensitivity to Group 3 fungicides (DMI) and reduced sensitivity to Group 7 (SDHI) have been detected in WA.

Barley spot-form net blotch: Resistance and reduced sensitivity to Group 3 fungicides (DMI) and Group 7 SDHI (fluxapyroxad) have been detected in WA.

A way to reduce the resistance development of the pathogen is to use fungicide mixtures with different modes of action. For example, fungicides containing both a strobilurin (Group 11) and uncompromised triazole (Group 3) active ingredient should have uncompromised activity.

Use an integrated disease management strategy, including growing a variety with a good disease resistance profile, managing regrowth barley, using crop rotation, and avoiding growing extremely dense canopy crops. As a part of the fungicide resistance management strategy, fungicides need to be used as a last resort. Refer to the department website at dpiird.wa.gov.au, and the Australian Fungicide Resistance Extension Network (AFREN) website at afren.com.au, for more information on barley net blotches, and powdery mildew.

Fungicide groups, mode of action: Group 3: Azoles/demethylase inhibitors (DMIs), Group 7: Succinate dehydrogenase inhibitors (SDHIs), Group 11: Strobilurins/quinone outside inhibitors (Qols), Group 5: Amines/Morpholines, Group 13: Azanaphthalene, more information available on the website of the Australian Fungicide Resistance Extension Network (AFREN) afren.com.au.

Contacts

For more information please contact:

- **Andrea Hills**, Email andrea.hills@dpird.wa.gov.au, Phone 08 9083 1144
- **Kith Jayasena**, Email Kithsiri.jayasena@dpird.wa.gov.au, Phone 08 9892 8477
- **Geoff Thomas**, Email Geoff.j.thomas@dpird.wa.gov.au, Phone 08 9368 3262
- **Ciara Beard**, Email ciara.beard@dpird.wa.gov.au, Phone 08 9956 8504

Please inform Ciara Beard of any errors or omissions. Email ciara.beard@dpird.wa.gov.au.

Last updated November 2024

Important Disclaimer

Mention of trade names does not imply endorsement or preference of any company's product by Department of Primary Industries and Regional Development (DPIRD) and any omission of a trade name is unintentional. Recommendations are current at the time this page was prepared.

The Chief Executive Officer of the Department of Primary Industries and Regional Development and the State of Western Australia accept no liability whatsoever by reason of negligence or otherwise arising from the use or release of this information or any part of it.

Copyright © State of Western Australia (Department of Primary Industries and Regional Development) 2024.

dpird.wa.gov.au